



**IV Iron Replacement Therapy for Iron Deficiency Anemia  
in Pregnancy (2<sup>nd</sup> & 3<sup>rd</sup> Trimester)  
– OUTPATIENT ORDERS**

**General**

☒ ALLERGY: \_\_\_\_\_

Diagnosis (check all that apply):

☐ O99.012 Anemia complicating pregnancy, second trimester

☐ O99.013 Anemia complicating pregnancy, third trimester

☐ D50.9 Iron deficiency anemia, unspecified

☐ D50.8 Other Iron Deficiency Anemias

☒ Metric Measurements: Weight: \_\_\_\_\_ kilograms; Height: \_\_\_\_\_ centimeters.

☒ OK to start infusion on date \_\_\_\_\_ based on labs dated \_\_\_\_\_

☒ Number of weeks gestation at time of order \_\_\_\_\_

**Laboratory**

☒ Ensure ordering provider to include/ attach a copy of current (within 30 days) lab results with this order set:  
CBC, Iron Panel (Iron, Transferrin, % TRF) and Ferritin

**Nursing Orders**

☒ Hypersensitivity / Anaphylaxis Management for Outpatient Infusions / Desensitization Order-Set

\*Note: epinephrine and methylprednisolone (Solu-medrol) are pregnancy category C\*

☒ Vital signs: baseline and every 30 minutes until discharge

☒ Hold patient 30 minutes post-infusion to observe for any signs and symptoms of hypersensitivity reactions

**Hydration Orders**

☐ Sodium chloride 0.9% 500 mL IV x 1 bag at 500 mL/hour as needed to decrease vein discomfort

**Pre-Medications**

☐ Acetaminophen (Tylenol) 650 mg orally x 1 dose (Pregnancy Category C)

☐ Diphenhydramine (Benadryl) 25 mg ☐ orally ☐ IV x 1 dose (Pregnancy category B)

☐ Other \_\_\_\_\_

**Medication Orders**

**Iron Sucrose (Venofer)**

☐ Iron sucrose 200 mg in sodium chloride 0.9% 100 mL IV over 30 minutes once a week x 5 doses

☐ Iron sucrose 200 mg in sodium chloride 0.9% 100 mL IV over 30 minutes twice a week x 5 doses

☐ Other \_\_\_\_\_

MD Print name \_\_\_\_\_

MD Contact #: \_\_\_\_\_

MD Signature \_\_\_\_\_

DATE \_\_\_\_/\_\_\_\_/\_\_\_\_ TIME \_\_\_\_:\_\_\_\_

Patient Name:

Date of Birth: