



General

☒ ALLERGY: _____

PRIMARY DIAGNOSIS: *** Must choose a primary from lists below ***

- ☐ D50.0 Iron deficiency anemia secondary to blood loss (chronic) ☐ D64.81 Anemia due to antineoplastic chemotherapy
☐ D50.8 Other Iron Deficiency Anemias ☐ D63.1 Anemia in chronic kidney disease

SECONDARY DIAGNOSIS:

- ☐ K90.9 Intestinal malabsorption, unspecified ☐ N18.9 Chronic kidney disease, unspecified
☐ T50.905A Unspecified adverse effect of unspecified drug, medicinal and biological substance
☐ T50.995A Other drug allergy

☒ Metric Measurements: Weight: _____ kilograms; Height: _____ centimeters.

☒ OK to start infusion on date _____ based on labs dated _____

Laboratory

- ☒ Ensure ordering provider to include/ attach a copy of current (within 30 days old) lab results (baseline CBC, Iron Panel (Iron, Transferrin, % TRF, and Ferritin) with this order-set

Nursing Orders

- ☒ IV ACCESS: ☐ Peripheral ☐ PICC ☐ Tunneled Catheter ☐ Implanted Port ☐ Other _____
☐ May Access Port-A-Cath ☐ May draw labs from Central Line
- ☒ Hypersensitivity / Anaphylaxis Management for Outpatient Infusions / Desensitizations Order-Set
- ☒ Vital signs: baseline and every 30 minutes until discharge
- ☒ Hold patient 30 minutes post-infusion to observe for any signs and symptoms of hypersensitivity reactions

Hydration Orders

- ☒ Sodium chloride 0.9% 500 mL IV x 1 bag at 500 mL/hour as needed to decrease vein discomfort

Medication Orders

Ferric Derisomaltose (Monoferric)

- ☐ Ferric Derisomaltose 1000mg in sodium chloride 0.9% 250 mL IV over 20 minutes x 1 dose. If Monoferric is not the preferred agent, may use (Venofer) Iron sucrose 300 mg in sodium chloride 0.9% 250 mL IV over 90 minutes every 2 days x 3 doses
If weight less than 50 kg, pharmacist to calculate dose at 20 mg/kg of body weight

Ferric Carboxymaltose (Injectafer)

- ☐ Ferric carboxymaltose 750 mg in sodium chloride 0.9% 250 mL IV over 30 minutes x 1 dose
☐ Ferric carboxymaltose 750 mg in sodium chloride 0.9% 250 mL IV over 30 minutes every 7 days x 2 doses
If weight less than 50 kg, pharmacist to notify MD for dose calculation at 15 mg/kg of body weight

Ferumoxytol (Feraheme) for chronic kidney disease, unspecified (D63.1/N18.9) ONLY

- ☐ Ferumoxytol 510 mg in sodium chloride 0.9% 100 mL IV over 30 minutes x 1 dose
☐ Ferumoxytol 510 mg in sodium chloride 0.9% 100 mL IV over 30 minutes every 7 days x 2 doses

Iron Sucrose (Venofer)

- ☐ Iron sucrose 300 mg in sodium chloride 0.9% 250 mL IV over 90 minutes every 2 days x 3 doses

MD Print name _____

MD Contact #: _____

MD Signature _____ DATE ____/____/____ TIME ____:____

Reviewed: 09/2026

Patient Name:

Date of Birth: